

AD _____

Award Number: DAMD17-01-1-0199

TITLE: Use of a Transgenic Mouse Model with a Regulatable Estrogen Receptor Alpha (ER) to Study the Role of ER in Mammary Gland Development and Cancer

PRINCIPAL INVESTIGATOR: Sonia L. Sugg, M.D.

CONTRACTING ORGANIZATION: Medical College of Wisconsin
Milwaukee, WI 53226-0509

REPORT DATE: September 2003

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20040329 025

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 074-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE September 2003		3. REPORT TYPE AND DATES COVERED Annual (20 Aug 2002 - 19 Aug 2003)
4. TITLE AND SUBTITLE Use of a Transgenic Mouse Model with a Regulatable Estrogen Receptor Alpha (ER) to Study the Role of ER in Mammary Gland Development and Cancer			5. FUNDING NUMBERS DAMD17-01-1-0199	
6. AUTHOR(S) Sonia L. Sugg, M.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Medical College of Wisconsin Milwaukee, WI 53226-0509 E-Mail: ssugg@mcw.edu			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Original contains color plates: ALL DTIC reproductions will be in black and white				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 Words) Estrogen (E2) is required for the development of breast cancer, but there are few animal models to study its mechanism of action in vivo. We hypothesize that the timing of estrogen receptor α (ER α) action is crucial in normal mammary gland development and tumorigenesis. Therefore, a transgenic mouse model with a regulatable ER α is being generated. In vitro, a mutant ER α (525L) has a severely attenuated response to endogenous estradiol (E2) but a wild-type (WT) response to diethylstilbestrol (DES). Gene targeting was used to insert 525L into the WT ER gene. Positive clones were injected into mouse blastocysts and chimeras were generated. There has been no germline transmission so far. Additional injections and further breeding is in progress. The resultant transgenic mouse will enable us to study how the timing of ER activation modulates normal mammary maturation and the development of mammary cancers.				
14. SUBJECT TERMS Breast cancer, estrogen receptor			15. NUMBER OF PAGES 7	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

Table of Contents

Cover letter	Page 1
SF 298	Page 2
Introduction	Page 4
Body	Page 4
Key Research Accomplishments.....	Page 6
Reportable Outcomes	Page 7
Conclusion	Page 7
References.....	Page 7

INTRODUCTION

Estrogen and its receptor are required for normal breast development and are important in the treatment and prevention of breast cancer. However, there are few suitable animal models to study the role of ER activation in the breast development and cancer. We developed a mutant ER having a severely attenuated response to endogenous estradiol while demonstrating a wild-type response to a synthetic estrogen agonist *in vitro*. This was due to a difference in binding affinity. We postulate that the timing and nature of estrogen action is crucial in normal mammary gland development and tumorigenesis. We are proposing a transgenic mouse model with a regulatable ER to test this hypothesis. Our specific aims are to 1) Develop transgenic mice using gene-targeting techniques expressing only the ligand discriminating mutant ER. 2) Define the contribution of ligand induced transcriptional activation of ER in the development of normal mammary gland. 3) Investigate how the timing of ER activation modulates the development of genetically and chemically induced mammary cancers. The mutant ER mice are uniquely suited to study the role of ligand independent ER activation *in vivo*, and will also be useful in studying reproductive tract development and behavior.

BODY

Task 1. Transgenic mice expressing only a mutant ER with ligand discrimination will be developed. (months 1 - 12)

a) Mouse ER genomic DNA will be cloned. A targeting construct (8 to 10 kb) will be generated.
Completed.

b) Mice transgenic and homozygous for mutant ER will be generated.
A two-pronged effort, with 2 different targeting constructs, was started at MCW and the University of Chicago. The rationale was to increase the probability of, and decrease the time required to successfully generating a transgenic mouse.

1) Work performed at the University of Chicago.

An 11.5 kilobase ER α construct containing the G525L mutation (purple bar in green exon 9 box), an 18 bp 6xHis-tag epitope, and an extra Xba1 site, was engineered to facilitate homologous recombination into the mouse genome (Figure 1). The original construct was obtained from Dr. Kenneth Korach¹ and the G525L mutation was inserted in our laboratory by site directed mutagenesis. We also inserted an ACN cassette into the targeting construct, obtained from Dr. Wondisford at the University of Chicago, which contained a testis-specific




Fig. 1. ER α construct containing the G525L mutation with ACN cassette

promoter (tACE), Cre structural gene, mouse RNA polymerase gene, and neo cassette, flanked at the 5' and 3' ends by loxP sites² (Figure 1).

The targeting construct was electroporated into ES cells. Positive clones were identified by 5' external probe Southern blots (Figure 2). A PCR strategy to identify 3' integration was also developed (Figure 3). PCR of exon 9 was used to detect the 6xHis-tag, Xba1 site, and G525L mutation in the positive clones (Figure 4). Southern blots with an internal probe were also performed to demonstrate only one insertion event in each positive clone (Figure 2). Two positive clones, 55 and 71, were obtained. They were injected into mouse blastocysts and four chimeras, one female from clone 55 and two males and one female from clone 71, were generated. Thus far, germline transmission has not been achieved with any of the chimeras.

In the future, electroporation of the targeting construct into fresh ES cells will be repeated. With the assistance of our collaborators, who have extensive expertise in developing knock-in and knock-out mouse models, we are confident that this approach will be successful. If not, an alternate strategy of inserting the mutant ER α into an α ERKO background could be used to create a similar model. In this system, ER α expression would be driven by an exogenous promoter.




Fig. 2 Identification of positive clones by 5' external probe Southern blots




Fig. 3. PCR strategy to identify 3' integration




Fig. 4. PCR detection of G525L mutant in positive ES clones

After germline transmission, Southern blot genotyping from tail DNA restriction digest will confirm the excision of the ACN cassette. The tACE promoter will drive expression of the Cre-recombinase gene in the testis of the chimeric mice during spermatogenesis, so that the ACN cassette is self-excised in the ES cell derived sperm of the chimeras². A Southern blot strategy to detect the ACN cassette excision in the F1 generation has been developed (Figure 5).




Fig. 5 Detection of ACN

2) Work performed at the Medical College of Wisconsin.

A 15 kb G525L mutant ER targeting construct was used to generate 18/380 (4.7%) positive ES cell clones, as detected by 5' probe Southern blot screening (for details please see last year's report). The positive clones were further confirmed by 3' external probe Southern hybridization¹, PCR of exon 9 followed by restriction enzyme XbaI digest (Figure 4), G525L mutation sequencing, and PCR of the neo selection

Table 1					
ES Clones#	Southern/5'probe	Seq.PCR/XbaI	Southern/3'probe	Mutation Sequencing	Southern/Neo
#362	+++++		+++	+	+
#29	+++++		+++	+	+
#2	++++		-	N/A	+
#200	++++		+++	+	+
#110	+++		++	+	+
#130	+++		-	+	+
#134	+++	non cut	N/A	N/A	+
#175	+++		-	N/A	+
#241	+++		+	+	+
#244	+++		+++	+	+
#269	+++		-	N/A	+
#299	+++		+++	+	+
#28	++	non cut	N/A	N/A	+
#34	++		-	N/A	+
#225	++	non cut	N/A	N/A	+
#257	++		++	+	+
#331	++	non cut	N/A	N/A	+
#338	++	non cut	N/A	N/A	+
#1	-		-	-	-
#3	-	cut	+++	+	+
#4	-		-	N/A	-
#5	-		-	-	-
#6	-	cut	++	+	-
#13	-		-	N/A	-
#14	-		-	-	-
#16	-		-	N/A?	-
#17	-		++	N/A	+
#18	-		-	N/A	-

marker (see Table 1). Three positive clones 29, 200, and 362 were selected for blastocyst injection. The genomic DNA from these 3 clones was again confirmed by all of the above methods to be positive prior to blastocyst injection. Clones 200 and 362 had litters that were aborted or cannibalized. Clone 29 generated 6 chimeras with 60 to 90% coat color chimerism. The first breeding generated no germline transmission. We are continuing to breed clone 29 chimeras and are reinjecting clones 200 and 362.

KEY RESEARCH ACCOMPLISHMENTS:

- 1) Generation of mutant ER chimeras from 2 targeting constructs.

REPORTABLE OUTCOMES:

Abstracts:

- 1) Sugg SL, Li J, Huang JJ, Radek J, Brace J, Swope DL, Korach KS, Greene GL. Design of a transgenic mouse with a regulatable estrogen receptor alpha. Era of Hope Department of Defense Breast Cancer Research Program Meeting, Orlando 2002.
- 2) Sugg SL, Li J, Radek J, Shilyansky J, Greene GL. Differences in Binding Affinity Causes Mutant Estrogen Receptor Alpha Ligand Discrimination. Proc AACR, 2003.

CONCLUSIONS

Over the second year of this award, we have made substantial progress toward creating the transgenic mouse model with a regulatable ER that was proposed for this project. We anticipate having transgenic animals imminently. We will then be able to characterize the phenotype, followed by experiments to determine how the timing of ER action affects mammary gland maturation and modulates mammary tumorigenesis and latency in models of genetically and chemically induced mammary cancers. With the widespread clinical use of ER ligands, detailed understanding of ER action is increasingly important. This model will allow us to manipulate ER transcriptional activity *in-vivo* during different developmental stages, and will add significantly to current knowledge on ligand-activated and ligand-independent ER action. Results from our proposal on how the timing of ER activation modulates normal mammary maturation and the development of mammary cancers will likely lead to determining susceptible periods for treatment in the prevention of breast cancer in humans.

REFERENCES

1. Swope DL, Castranio T, Harrell JC, Mishina Y, Korach KS. AF-2 knock-in mutation of estrogen receptor α : cre-loxP excision of a PGK-neo cassette from the 3' UTR. *Genesis*. 2002;32:99-101.
2. Bunting M, Bernstein KE, Greer JM, Capecchi MR, Thomas KR. Targeting genes for self-excision in the germ line. *Genes & Development*. 1999;13:1524-8.